

**RESEARCH PROBLEMS
FROM THE 18TH WORKSHOP ‘3IN1’ 2009**

edited by Mariusz Meszka

Abstract. A collection of open problems that were posed at the 18th Workshop ‘3in1’, held on November 26-28, 2009 in Krakow, Poland. The problems are presented by Zdenek Ryjacek in “Does the Thomassen’s conjecture imply $N=NP$?” and “Dominating cycles and hamiltonian prisms”, and by Carol T. Zamfirescu in “Two problems on bihomogeneously traceable digraphs”.

Keywords: Hamilton-connected graph, hamiltonian graph, dominating cycle, bihomogeneously traceable graph.

Mathematics Subject Classification: 05C45, 68Q25, 05C38, 05C20.

1. DOES THE THOMASSEN’S CONJECTURE IMPLY $N=NP$?

Zdeněk Ryjáček
ryjacek@kma.zcu.cz
University of West Bohemia
Department of Mathematics
Czech Republic

By a *graph* we mean a simple loopless finite undirected graph $G = (V(G), E(G))$. A graph G is *Hamilton-connected* if G has a hamiltonian (x, y) -path for any $x, y \in V(G)$, and, for an integer $k \geq 1$, G is *k -Hamilton-connected* if $G - X$ is Hamilton-connected for any $X \subset V(G)$ with $|X| = k$. Denote $E^+(G) = \{xy \mid x, y \in V(G)\}$, and for $X \subset E^+(G)$ set $G + X = (V(G), E(G) \cup X)$ (i.e., X is a set of “new” edges that are “added” to G ; if $e_1 = \{x, y\} \in E(G)$ and $e_2 = \{x, y\} \in X$, we consider e_1 and e_2 as parallel edges of $G + X$). A graph G is said to be *k -edge-Hamilton-connected* if, for any $X \subset E^+(G)$ such that $|X| = k$ and the edges of X determine a path system, the graph $G + X$ has a hamiltonian cycle containing all edges in X . The following facts are easy to observe.

1. A graph G is 1-edge-Hamilton-connected if and only if G is Hamilton-connected.

2. A graph G is 2-edge-Hamilton-connected if and only if:
- (i) G is 1-Hamilton-connected (i.e., $G - x$ is Hamilton-connected for any vertex $x \in V(G)$), and
 - (ii) for any four distinct vertices $x_1, x_2, x_3, x_4 \in V(G)$, G has a path factor consisting of 2 paths P_1, P_2 such that both P_1 and P_2 have one endvertex in $\{x_1, x_2\}$ and one endvertex in $\{x_3, x_4\}$.
3. If G is 2-edge-Hamilton-connected, then G is 4-connected.

Consider the following two decision problems.

k -E-HC

Instance: A graph G .

Question: Is G k -edge-Hamilton-connected?

k -E-HCL

Instance: A line graph G .

Question: Is G k -edge-Hamilton-connected?

(i.e., k -E-HCL is k -E-HC restricted to line graphs).

Question 1: Determine the complexity of 2-E-HCL.

The following facts are known:

- **HAM**
Instance: A graph G .
Question: Does G contain a hamiltonian cycle?
 HAM \in NPC, even if restricted to line graphs.
- **H-PATH**
Instance: A graph G and distinct vertices $u, v \in V(G)$.
Question: Does G contain a hamiltonian (u, v) -path?
 H-PATH \in NPC, even if restricted to line graphs [1].
- **H-CONN**
Instance: A graph G .
Question: Is G Hamilton-connected?
 H-CONN \in NPC [3].
- **1-H-CONN**
Instance: A graph G .
Question: Is G 1-Hamilton-connected?
 1-H-CONN \in NPC [6].

Thus, a common guess would be that probably 2-E-HCL \in NPC.

Question 2: Why is Question 1 interesting?

The following conjecture was posed in [5].

Conjecture [Thomassen]. Every 4-connected line graph is hamiltonian.

There are many known equivalent versions of the Thomassen's conjecture; among others, we mention the following.

Theorem. The following statements are equivalent:

- (i) Every 4-connected line graph is hamiltonian.

- (ii) Every 4-connected line graph is 2-edge-Hamilton-connected [4].
- (iii) Every snark has a dominating cycle [2].

Thus, if the Thomassen’s conjecture is true, then a line graph G is 2-edge-Hamilton-connected if and only if G is 4-connected, implying that 2-E-HCL is polynomial. Consequently, proving the “common guess” $2\text{-E-HCL} \in \text{NPC}$ would mean

- disproving the Thomassen’s conjecture,
- proving the existence of a snark with no dominating cycle,

unless $P=NP$.

REFERENCES

- [1] A.A. Bertossi, *The edge hamiltonian path problem is NP-complete*, Inform. Process. Lett. **13** (1981), 157–159.
- [2] H.J. Broersma, G. Fijavž, T. Kaiser, R. Kužel, Z. Ryjáček, P. Vrána: *Contractible subgraphs, Thomassen’s conjecture and the dominating cycle conjecture for snarks*, Discrete Mathematics **308** (2008), 6064–6077.
- [3] Alice M. Dean, *The computational complexity of deciding Hamiltonian-connectedness*, Proceedings of the Twenty-fourth Southeastern International Conference on Combinatorics, Graph Theory, and Computing (Boca Raton, FL, 1993). Congr. Numer. **93** (1993), 209–214.
- [4] R. Kužel, Z. Ryjáček, P. Vrána, *On some conjectures on hamiltonian properties of 4-connected line graphs*, Preprint 2009, to appear.
- [5] C. Thomassen, *Reflections on graph theory*, J. Graph Theory **10** (1986), 309–324.
- [6] P. Vrána, personal communication.

2. DOMINATING CYCLES AND HAMILTONIAN PRISMS

Zdeněk Ryjáček
 ryjacek@kma.zcu.cz
 University of West Bohemia
 Department of Mathematics
 Czech Republic

The *prism over a graph G* , denoted $G \square K_2$, is the Cartesian product of G and K_2 . It consists of two disjoint copies of G and a perfect matching connecting a vertex in one copy of G to its “clone” in the other copy.

A graph G is *hamiltonian* if it has a hamiltonian cycle and *traceable* if it has a hamiltonian path. Define a *k-walk* in a graph to be a spanning closed walk in which every vertex is visited at most k times

The following implications are easy to verify:

G is hamiltonian $\Rightarrow G$ is traceable $\Rightarrow G \square K_2$ is hamiltonian $\Rightarrow G$ has a 2-walk.


Thus the question whether G has a hamiltonian prism (i.e whether $G \square K_2$ is hamiltonian) is “sandwiched” between hamiltonicity and having a 2-walk. Specifically, the property of having a hamiltonian prism can be considered as a “relaxation” of hamiltonicity. More information about prism-hamiltonicity of a graph can be found e.g. in [1] and [2].

A *dominating cycle* in a graph G is a cycle C such that every edge of G has at least one vertex on C , i.e. such that the graph $G - C$ is edgeless. Clearly, a hamiltonian cycle is dominating, and hence the property of having a dominating cycle can be considered as another relaxation of hamiltonicity.

There is a natural question whether there is any relation between these two properties.

Example 1. Let H be any 2-connected cubic nonhamiltonian graph, and let G be obtained from H by replacing every vertex of H with a triangle (such a G is sometimes called the *inflation* of H). Then G is a 2-connected line graph and these are known [2] to be prism-hamiltonian. On the other hand, since H is nonhamiltonian, any cycle in G has to miss at least one “new” triangle and hence G has no dominating cycle. Thus, there are “many” graphs showing that hamiltonian prism does not imply having a dominating cycle.

Example 2. The graph in the figure below shows that also the existence of a dominating cycle does not imply having hamiltonian prism.



However, all such known examples are of low toughness (recall that G is 1-tough if, for any $S \subset V(G)$, the graph $G - S$ has at most $|S|$ components). This motivates the following question.

Conjecture. *Let G be a 1-tough graph having a dominating cycle. Then G has hamiltonian prism.*

Comments. Suppose that G has a dominating cycle C of even length. Set $M = V(G) \setminus V(C)$ and $N = \{x \in V(C) \mid x \text{ has a neighbor in } M\}$. Then the graph induced by $M \cup N$ has a matching containing all vertices from M (this follows by the toughness assumption and by the Hall’s theorem). Using this matching, it is easy to construct a hamiltonian cycle in $G \square K_2$.

The difficult case is when all dominating cycles in G are of odd length.

REFERENCES

- [1] R. Čada, T. Kaiser, M. Rosenfeld, Z. Ryjáček, *Hamiltonian decompositions of prisms over cubic graphs*, Discrete Mathematics **286** (2004), 45–56.
- [2] T. Kaiser, D. Král, M. Rosenfeld, Z. Ryjáček, H.-J. Voss, *Hamiltonian cycles in prisms over graphs*, Journal of Graph Theory **56** (2007), 249–269.

3. TWO PROBLEMS ON BIHOMOGENEOUSLY TRACEABLE DIGRAPHS

Carol T. Zamfirescu
 czamfirescu@gmail.com
 Technische Universität Dortmund
 Vogelpothsweg 87, 44221 Dortmund
 Germany

We concern ourselves here exclusively with simple finite oriented graphs (i.e. digraphs with no multiple edges, a finite number of vertices, and without cycles of length 2), calling these simply *graphs*. A graph is called *homogeneously traceable*, if for every vertex v there exists a hamiltonian path starting at v . If, additionally, the graph has the property that in every vertex a hamiltonian path ends, we call it *bihomogeneously traceable*. In this setting, and in a graph on n vertices, *arc-minimality* (or 2-regularity) means that the graph has precisely $2n$ edges (i.e. every vertex has in-degree 2 and out-degree 2). We remark that bihomogeneous traceability does not imply hamiltonicity, for instance hypohamiltonian graphs are non-hamiltonian and bihomogeneously traceable.

Z. Skupień [3] presented in 1981 an infinite family of arc-minimal non-hamiltonian bihomogeneously traceable graphs, featuring graphs of all orders greater or equal to 7. Another such infinite family of graphs (but not arc-minimal) was provided independently by S. Hahn and T. Zamfirescu [1] in the same year.

In 1983, L. E. Penn and D. Witte [2] proved that the cartesian product of two oriented cycles of length a and b is hypohamiltonian (whence, non-hamiltonian and bihomogeneously traceable) if and only if there exist relatively prime numbers $m, n \in \mathbb{N}$ such that $am + nb = ab - 1$. We note that these graphs are also arc-minimal.

In their 1981 paper, Hahn and Zamfirescu presented two planar non-hamiltonian bihomogeneously traceable graphs, one of which is arc-minimal, and asked the natural question whether infinitely many such graphs do exist. Very recently it was proven that this is indeed the case, see [4].

The following problems, however, are still open.

Problem 1. Is there an infinite family of planar arc-minimal non-hamiltonian bihomogeneously traceable oriented graphs?

Problem 2. Are there such graphs on all orders greater than some integer? Even if one removes the condition of arc-minimality, this problem is still open.

REFERENCES

- [1] S. Hahn, T. Zamfirescu, *Bihomogeneously traceable oriented graphs*, Rend. Sem. Mat., Torino **39** (1981) 2, 137–145.
- [2] L.E. Penn, D. Witte, *When the Cartesian product of two directed cycles is hypo-Hamiltonian*, J. Graph Theory **7** (1983), 441–443.
- [3] Z. Skupień, *On homogeneously traceable nonhamiltonian digraphs and oriented graphs*, Proc. Fourth Int. Conf. on the Theory and Appl. of Graphs, Kalamazoo (MI) 1980, Wiley (1981), 517–527.
- [4] C.T. Zamfirescu, *An infinite family of planar non-hamiltonian bihomogeneously traceable oriented graphs*, to appear in Graphs and Combinatorics.

Mariusz Meszka
meszka@agh.edu.pl

AGH University of Science and Technology
Faculty of Applied Mathematics
al. Mickiewicza 30, 30-059 Cracow, Poland

Received: November 28, 2009.

